

**Stellar Atmospheres:
Structure, Composition
and
Limb Darkening**

*H. M. Dyck
US Naval Observatory
Flagstaff, Arizona*

2000 MICHELSON SUMMER SCHOOL

References

○ Some general texts

Chandrasekhar, S. 1960, *Radiative Transfer* (Dover: New York).

Collins, G. W. 1989, *Fundamentals of Stellar Astrophysics* (W. H. Freeman: New York).

Shu, F. H. 1991, *The Physics of Astrophysics. Volume I. Radiation* (Univ. Science Books: Mill Valley, CA).

○ Selected papers

Bessell, M. S. *et al.* 1989, *A&A*, 213, 209.

Burns, D. *et al.* 1997, *MNRAS*, 290, L11.

Buscher, D. F. *et al.* 1990, *MNRAS*, 245, 7.

Claret, A. 1998, *A&AS*, 131, 395.

2000 MICHELSON SUMMER SCHOOL

Claret, A. *et al.* 1995, *A&AS*, 114, 247.

Gilliland, R. L. & Dupree, A. K. 1996, *ApJ*, 463, 29.

Hajian, A. R. *et al.* 1998, *ApJ*, 496, 484.

Hanbury Brown, R. *et al.* 1974, *MNRAS*, 167, 475.

Hestroffer, D. 1997, *A&A*, 327, 199.

Hofmann, K.-H. & Scholz, M. 1998, *A&A*, 335, 637.

Jacob, A. P. *et al.* 2000, *MNRAS*, 312, 733.

Kurucz, R. L. 1979, *ApJS*, 40, 1.

Manduca, A. *et al.* 1977, *A&A*, 61, 809.

Michelson, A. A. & Pease, F. G. 1921, *ApJ*, 53, 249.

Petro, L. D. *et al.* 1984, *ApJ*, 283, 426.

2000 MICHELSON SUMMER SCHOOL

Pierce, A. K. & Waddell, J. H. 1961, *MemRAS*, 58, 89.

Quirrenbach, A. *et al.* 1996, *A&A*, 312, 160.

Scholz, M. 1985, *A&A*, 145, 251.

Scholz, M. 2000, preprint.

Scholz, M. & Takeda, Y. 1987, *A&A*, 186, 200.

Tsuji, T. 1978, *PASJ*, 30, 435.

van Hamme, W. 1993, *AJ*, 106, 2096.

Weiner, J. *et al.* 2000, *ApJ*, in press.

White, N. M. 1980, *ApJ*, 242, 646.

Wilson, R. W. *et al.* 1992, *MNRAS*, 257, 369.


2000 MICHELSON SUMMER SCHOOL

Introductory remarks:

- **Resolved stars (*e.g.* the Sun) are observed to be darker at the edge (*i.e.* limb) than at the center of the disk.**
- **The phenomenon arises because stars have a temperature gradient — they are hotter in the deeper parts than in the outer parts of the atmosphere.**
- **By studying limb darkening at a variety of wavelengths we may hope to understand something about the atmospheric chemical composition and structure.**
- **Limb darkening studies, once restricted to very few stars, now broadly possible with the advent of optical interferometers.**

2000 MICHELSON SUMMER SCHOOL

Limb-darkening geometry:




Definitions:

- R = radius of the star.
- r = distance projected along disk from center.
- θ = angle between the normal to the stellar surface and the line of sight to the observer.
- $\mu = \cos \theta$.

2000 MICHELSON SUMMER SCHOOL

Simplified picture of limb darkening:




2000 MICHELSON SUMMER SCHOOL

Our expectations:

- The observed intensity is a function of T, τ_λ :

$$I_\lambda(\mu) = f[B_\lambda(T), \tau_\lambda]$$

- Expect lower intensities at limb than at center:



- Observe $I_\lambda(\mu)$ and infer T, τ_λ with the aid of radiative transfer models

2000 MICHELSON SUMMER SCHOOL

Radiative transfer formalism:

- The emergent intensity in the plane-parallel approximation is

$$I_{\lambda}(\mu) = \int_0^{\infty} S_{\lambda}(\tau_{\lambda}) e^{-\tau_{\lambda}/\mu} d\tau_{\lambda}/\mu$$

where $S_{\lambda}(\tau_{\lambda})$ is the *source function*.

- $S_{\lambda}(\tau_{\lambda})$, τ_{λ} carry information about chemical composition, temperature and pressure in the stellar atmosphere.

2000 MICHELSON SUMMER SCHOOL

How to make some progress:

- The relationship between intensity and source function is a Laplace transform:

$$\mu I_{\lambda}(\mu) = \mathcal{L}[S_{\lambda}(\tau_{\lambda})]$$

- Observe $I_{\lambda}(\mu)$, find the inverse Laplace transform and obtain $S_{\lambda}(\tau_{\lambda})$.
- Assume LTE and obtain $T(\tau_{\lambda})$ because $S_{\lambda}(\tau_{\lambda}) = B_{\lambda}(\tau_{\lambda})$, the *Planck function*.

2000 MICHELSON SUMMER SCHOOL

The Eddington approximation:

- **Emergent intensity found to be**

$$I(\mu) = I(1) \frac{3}{5} \left(\mu + \frac{2}{3} \right)$$

- **Leads to the following source function**

$$S(\tau) = \frac{3}{4\pi} F \left(\tau + \frac{2}{3} \right)$$

where F is the flux of radiation.

- **Get relation between T and τ :**

$$T^4 = \frac{3}{4} T_{eff}^4 \left(\tau + \frac{2}{3} \right)$$

2000 MICHELSON SUMMER SCHOOL

Limb-darkening laws

- The limb-darkening law in the Eddington approximation is a *linear limb-darkening law*. A more general form is

$$I(\mu) = I(1)(a + b\mu)$$

and may be found in the papers by van Hamme (1993) and Claret *et al.* (1995).


- Other polynomial forms have been suggested.
- A useful general limb-darkening function first suggested by Michelson & Pease (1921) and recently advocated by Hestroffer (1997) is

$$I(\mu) = I(1)\mu^\alpha$$

2000 MICHELSON SUMMER SCHOOL

Solar limb-darkening:

- Petro *et al.* (1984) obtained the following mean Solar limb-darkening at a wavelength of 445.1 nm. The Eddington approximation to the limb-darkening is shown for comparison.




- Eddington slope limb-darkening law is a poor fit. Petro *et al.* use a 5th degree polynomial to fit the intensity, yielding S_λ represented by a 5th degree polynomial in τ_λ .

2000 MICHELSON SUMMER SCHOOL

Solar temperature distribution:

- The following plot compares the $T(\tau)$ derived from the Eddington approximation for the Sun to detailed model from Kurucz (1979).



- Eddington approximation temperatures agree with model to $\approx 2\%$ over the range $0.1 \leq \tau \leq 3$ even though limb-darkening fit is not good.
- Simple example illustrates how analysis of limb darkening leads to understanding of atmospheric temperature structure. Interpreting *real* observations is more complicated.


2000 MICHELSON SUMMER SCHOOL

Example inversion of real data:

- Solar limb-darkening data used by Pierce & Waddell (1961) to derive atmospheric temperature structure.
- Analysis of multi-wavelength data also shows how Solar continuous opacity derived from limb darkening observations.
- One of several investigations confirming H^- as an important constituent in the Solar atmosphere.
- Investigation shows how temperature *and* composition of the atmosphere may be determined from limb-darkening observations at a variety of wavelengths.

2000 MICHELSON SUMMER SCHOOL

Some observational data:



Notes:

- Horizontal row of dots at top are a uniformly-bright disk (*i.e.*, no limb darkening).
- Wavelengths for observed data are, starting from the lowest row of dots, 0.42, 0.6, 1.0 and 2.1 μm .

2000 MICHELSON SUMMER SCHOOL

Some comments:

- **Sharp edge to the Sun, evident at all wavelengths.**
- **Wavelength dependence to limb darkening with near infrared behaving most like a uniform disk.**

2000 MICHELSON SUMMER SCHOOL

What's next?

- Assume that the source function may be written as the following expansion:

$$\frac{S_{\lambda}(\tau_{\lambda})}{I_{\lambda}(1)} = a_{\lambda} + b_{\lambda} \tau_{\lambda} + c_{\lambda} E_2(\tau_{\lambda})$$

where $E_2(\tau_{\lambda})$ is an exponential integral and a_{λ} , b_{λ} and c_{λ} are constants to be determined.

- This results in a limb darkening function of the form:

$$\frac{I_{\lambda}(\mu)}{I_{\lambda}(1)} = a_{\lambda} + b_{\lambda} \mu + c_{\lambda} [1 - \mu \ln(1 + \mu^{-1})]$$

- Fit the observed limb darkening data and determine the value of the constants at each wavelength.

2000 MICHELSON SUMMER SCHOOL

- Assume LTE which gives

$$\frac{S_{\lambda}(\tau_{\lambda})}{I_{\lambda}(1)} = \frac{B_{\lambda}(\tau_{\lambda})}{I_{\lambda}(1)}$$


- Measure the absolute intensity at disk center, $I_{\lambda}(1)$ and numerically compute a solution.
- Results in a table $T_{\lambda}(\tau_{\lambda})$ for all wavelengths observed.
- For a fixed T_{λ} , τ_{λ} will vary because of the changing opacity since, by definition,

$$\tau_{\lambda} = \int_x^0 k_{\lambda} \rho dx$$

over some path length x , where k_{λ} is the absorption coefficient of the material and ρ is the density.

2000 MICHELSON SUMMER SCHOOL

Results of the Pierce & Waddell study:



Bottom curve is shape of opacity derived from observations, upper curve is theoretical opacity for H^- .

2000 MICHELSON SUMMER SCHOOL

Other stars:

- Sun is special case since it is a very resolved disk.
- For other stars, interferometers must be used to resolve the disks.
- Interferometers measure the *visibility* rather than the center-to-limb variation (CLV) directly.
- Recall the Michelson & Pease CLV:

$$I(r/R) = I(0) [1 - (r/R)^2]^{\alpha/2}$$



- Hestroffer shows that this leads to the following visibility as a function of spatial frequency, s :

$$V_v(s) = \Gamma(v + 1) \frac{|J_v(s)|}{(s/2)^v}$$

where $v = (\alpha/2) + 1$, J_v is the v -th Bessel function of the first kind and Γ is the gamma function.

2000 MICHELSON SUMMER SCHOOL

Application to the Sun:



2000 MICHELSON SUMMER SCHOOL

Salient points of the previous graphs:

- **Michelson & Pease model fits CLV observations pretty well.**
- **Not much difference between uniform disk and limb-darkened visibilities where s is smaller than the first null.**
- **Limb-darkening information contained in height of secondary maxima and positions of nulls.**
- **Limb-darkening effect is small, so high accuracy measurements are needed.**

2000 MICHELSON SUMMER SCHOOL


Summary for stars with $T > 4000$ K:

- **Atmosphere models pretty good.**
- **Observed limb-darkening agrees with models.**
- **Any wavelength should yield good quality effective temperatures.**
- **Temperature structure and continuous opacity may be derived with adequate wavelength coverage.**
- **Earliest example was by Hanbury Brown *et al.* (1974) for Sirius, using the intensity interferometer.**
- **First Michelson interferometry observations beyond the first null of the visibility function were reported by Burns *et al.* (1997).**
- **Some other Michelson interferometry examples follow.**

2000 MICHELSON SUMMER SCHOOL

The next two plots are for α Boo, a K2 III star:


The plot below is from Quirrenbach *et al.* (1996) at 550 nm, using the Mark III interferometer.



Two attempted fits of uniform disk curves to the data are shown in the figure above.

2000 MICHELSON SUMMER SCHOOL


The plot below is from a variety of observations made at 2.2 μm , using different interferometers.



Here, the uniform disk fits reasonably well.

2000 MICHELSON SUMMER SCHOOL


Limb-darkened diameter versus wavelength for α Boo from Quirrenbach et al. (1996):



Note that the limb-darkened diameters are the same at all wavelengths, within the errors. Corrections for two different model temperatures are shown. The best determined diameter is 21.0 ± 0.2 mas.

2000 MICHELSON SUMMER SCHOOL

The following plot is from Hajian *et al.* (1998) for α Cas, a K0 III star:



The model above was taken from stellar atmosphere CLV calculations by Kurucz.

2000 MICHELSON SUMMER SCHOOL


What about the cooler stars?

- **Opacities dominated by lines of molecular species.**
- **Atmospheres may be very extended (*e.g.* the Mira variables & supergiants).**
- **Often don't have a sharp 'edge' to the star.**
- **Radius may be very strong function of wavelength, owing to two factors:**
 - **Limb darkening effects**
 - **Significant change of physical depth of the principal radiating layers.**


2000 MICHELSON SUMMER SCHOOL

Pictorial differences

- Warmer stars with compact atmospheres:




- Cooler stars with extended atmospheres:



2000 MICHELSON SUMMER SCHOOL

Computed spectrum for a $T = 3500$ K, $L = 500 L_{\odot}$ star, taken from Scholz & Takeda (1987):



2000 MICHELSON SUMMER SCHOOL

Where is the stellar surface?


- Introduce new concept — the *Rosseland mean radius*, R_{ROSS} — a fictitious surface:
 - Rosseland mean opacity is wavelength averaged opacity.

$$\frac{1}{k_R} = \frac{\int_0^\infty \frac{1}{k_\lambda} \frac{dB_\lambda(T)}{dT} d\lambda}{\int_0^\infty \frac{dB_\lambda(T)}{dT} d\lambda}$$

- R_{ROSS} defined as physical depth where $\tau_R = 1$.
- Used in effective temperature calculations.
- Must relate observed visibilities to this surface through models.
- Visibilities may not have same character as visibilities for warmer stars.

2000 MICHELSON SUMMER SCHOOL


Model CLV for the $T = 3500$ K, $L = 500 L_{\odot}$ model,
taken from Hofmann & Scholz (1998):



Note the fuzzy edge of the star, as seen in different wavelengths.


2000 MICHELSON SUMMER SCHOOL

Visibility functions for the same model:



2000 MICHELSON SUMMER SCHOOL


Model CLV for the $T = 3500$ K, $L = 10,000 L_{\odot}$ model, taken from Hofmann & Scholz (1998):



Note *VERY* fuzzy edge to the star for this model.


2000 MICHELSON SUMMER SCHOOL

Change of radius for the same model, taken from Scholz (1985):



2000 MICHELSON SUMMER SCHOOL

Visibility functions for the same model:




2000 MICHELSON SUMMER SCHOOL

Examples of observations of α Ori (M2Iab):

**Observations by Michelson & Pease (1921),
Weiner *et al.* (2000) and Gilliland & Dupree (1996) as
well as others summarized in White (1980) and Weiner
*et al.***

2000 MICHELSON SUMMER SCHOOL

The following is a plot of the uniform disk diameters for α Ori that are summarized in White (1980):




2000 MICHELSON SUMMER SCHOOL

Notes:

- Much of the scatter may be due to presence of TiO bands.
- Some scatter may result from time variability (White 1980).
- There is a suggestion of a decrease of diameter with increasing wavelength. Tsuji (1978) explained this by the addition of a circumstellar (scattering) dust shell to the photosphere. The effects of such shells is discussed by Scholz (2000).
- The presence of hot spots may complicate the interpretation of the diameter as a function of wavelength (Buscher *et al.* 1990, Wilson *et al.* 1992).

2000 MICHELSON SUMMER SCHOOL

In the following plot, the previous data plus data from Weiner *et al.* (2000) and Gilliland & Dupree (1996) are shown for α Ori. These extend from the UV to the IR.



2000 MICHELSON SUMMER SCHOOL

Notes:

- **The diameter appears to decrease from the UV into the red and then increase again toward the mid-IR.**
- **The larger diameter in the UV has been explained by Gilliland & Dupree (1996) as a measurement of the chromosphere rather than the photosphere.**
- **The slight increase in diameter from the near to the mid IR may be a contribution from the emission of the same dust postulated by Tsuji (1978). This needs to be tested.**

2000 MICHELSON SUMMER SCHOOL

Summary for cool, extended stars:

- **Variation of angular diameter with wavelength offers wealth of data against which to test spherical, extended model atmospheres (Scholz & Takeda 1987, Hofmann & Scholz 1998).**
- **Important sources of opacity may be characterized with the right kinds of observations (Jacob *et al.* 2000).**
- **Interpretation of visibility curves may be complicated by the existence of circumstellar shells or surface features.**